

Action-Angle Formulation of Charged Particle Dynamics in Toroidally Confined Plasmas

YANNIS KOMINIS

*School of Applied Mathematical and Physical Sciences
National Technical University of Athens, Athens 15780, Greece
gkomin@central.ntua.gr*

Charged particle dynamics under the presence of electromagnetic fields is one of the main paradigms where the modern theory of nonlinear Hamiltonian dynamics and chaos has been applied [1]. Such complex dynamics crucially determines particle energy and momentum transport in toroidally confined fusion plasmas. Particle dynamics are studied in terms of the *Guiding Center* (GC) approximation, and are integrable under magnetic field symmetry conditions, such as axisymmetry or helical symmetry. However, the presence of spontaneously or intentionally introduced symmetry-breaking, multi-scale electromagnetic perturbations results in non-integrable dynamics and chaos [2].

In this work, we introduce a novel integrable system for the unperturbed GC motion and we analytically calculate the *Action-Angle variables* of the system as well as the *Orbital Spectrum* consisting of the frequencies in all degrees of freedom [3]. The system serves as a starting point for the application of canonical perturbation theories and numerical studies under the presence of symmetry-breaking perturbations. The unique advantages of the Action-Angle formulation are discussed, in terms of time-scale separation, derivation of reduced kinetic models, a priori knowledge of the exact locations of the phase space where resonances take place, and their implications for charged particle transport [4].

References

- [1] A.J. Lichtenberg and M.A. Lieberman, *Regular and Chaotic Dynamics*, Springer (1992).
- [2] P.A. Zestanakis, Y. Kominis, G. Anastassiou, and K. Hizanidis, “Orbital spectrum analysis of non-axisymmetric perturbations of the guiding-center particle motion in axisymmetric equilibria”, *Phys. Plasmas* **23**, 032507 (2016).
- [3] Y. Antonenas, G. Anastassiou, and Y. Kominis, “Analytical calculation of the orbital spectrum of the guiding centre motion in axisymmetric magnetic fields”, *J. Plasma Phys.* **87**, 855870101 (2021).
- [4] Y. Kominis, A.K. Ram, and K. Hizanidis, “Kinetic Theory for Distribution Functions of Wave-Particle Interactions in Plasmas”, *Phys. Rev. Lett.* **104**, 235001 (2010).