

Existence of nonattainable states for Schrödinger type flows on the half space

Türker Özşarı

*Department of Mathematics
Bilkent University, Ankara 06800 Turkey
turker.ozsari@bilkent.edu.tr • <https://turkerozsari.com>*

Dedicated to 70th birthday of Professor Athanassios Fokas

In this talk, we address the boundary controllability problem for the Schrödinger equation on the half space (1D and 2D). We also extend the discussion to biharmonic Schrödinger equation. For both equations, we prove existence of unattainable states with L^2 type controls. Our method relies on recent developments in local wellposedness theory for initial boundary value problems for these equations ([1]-[4]). The noncontrollability results obtained here contrast with the dynamics in the case of bounded domains where controllability property holds true ([5]).

References

- [1] A. Batal, A. S. Fokas, and T. Özşarı, *Fokas method for linear boundary value problems involving mixed spatial derivatives*, Proc. A. **476** (2020), no. 2239, 20200076, 15.
- [2] Ahmet Batal and Türker Özşarı, *Nonlinear Schrödinger equations on the half-line with nonlinear boundary conditions*, Electron. J. Differential Equations (2016), Paper No. 222, 20.
- [3] Athanassios S. Fokas, A. Alexandrou Himonas, and Dionyssios Mantzavinos, *The nonlinear Schrödinger equation on the half-line*, Trans. Amer. Math. Soc. **369** (2017), no. 1, 681–709.
- [4] A. Alexandrou Himonas and Dionyssios Mantzavinos, *Well-posedness of the nonlinear Schrödinger equation on the half-plane*, Nonlinearity **33** (2020), no. 10, 5567–5609.
- [5] Elaine Machtyngier, *Exact controllability for the Schrödinger equation*, SIAM J. Control Optim. **32** (1994), no. 1, 24–34.